

Lysimachia loeflingii (arvensis) & foemina

2024-07-05

[İletişim \(FB\)](#) • [En yeni PDF](#) • [PDF'lar klasörü](#)

[Contact \(FB\)](#) • [Latest PDF](#) • [PDFs Folder](#)

Mavi Farekulagi / Blue Scarlet Pimpernel

Lysimachia loeflingii F.J.Jiménez & M.Talavera (= L. arvensis var. azurea = var. caerulea)

ile karşılaştırıldığında / compared with

Bagirsakotu / Blue Pimpernel

Lysimachia foemina (Mill.) U.Manns & Anderb.

Lysimachia loeflingii

Lysimachia foemina

Lysimachia loeflingii

Lysimachia foemina

Flowers

Yaprakları, geniş (neredeyse her zaman)
Güçlü değişmeyen mavi

Broad petals (nearly always)
Strong uniform blue

Yaprakları, daha dar (çok sık)
Mavinin farklı tonu (genellikle)
Çiçekler daha küçük olabilir



Narrower petals (very often)
Different tone of blue (usually)
Flowers may be smaller

Salgılı tüyleri, 3-hücreli (çok yayılmış)

Glands 3-celled (widely spread)

Salgılı tüyleri, (3)4-hücreli (genellikle daha az yayılmış)

Glanding (3)4-celled (often more restricted)




Fruits

5-6 damarlı

5-6 veined

Çok damarlı (neredeyse her zaman, gerekirse birkaçını kontrol edin)

Many-veined (virtually always, check several if needed)



Leaves

Geniş soluk yeşil (genellikle)
Çiçek sapları, uzun (genellikle)

*Broad pale-green leaves (usually)
Long flower stalks (usually)
(Upper leaves sometimes narrower,
and leaves sometimes darkish;
flower stalks a little variable)*

Daha dar daha koyu yeşil
Çiçek sapları, daha kısa (genellikle)

*Narrower darker green leaves
Shorter flower stalks (typically)*

İkişerli (neredeyse her zaman)

In 2s (almost always)

2 li (veya 3-lü)

In 2s (or 3s)



Plant



Gland Cells

3-hücreli tüyleri	4-hücreli tüyleri (bazen 3)		
3-celled glands	4-celled glands (sometimes 3)		
blue_celery @ iNaturalist	blue_celery @ iNaturalist		
